Taylor & Francis Online

Biotechnic & Histochemistry ›
Volume 92, 2017 - Issue 5

369 Views | 10 CrossRef citations to date | 1 Altmetric

Articles

Long-term exposure to a continuous 900 MHz electromagnetic field disrupts cerebellar morphology in young adult male rats

A Aslan, A İkinci, O Baş, OF Sönmez, H Kaya & E Odacı

Pages 324-330 | Accepted 20 Mar 2017, Published online: 16 May 2017

Cite this article https://doi.org/10.1080/10520295.2017.1310295 Check for updates

Sample our Medicine, Dentistry, Nursing & Allied Health Journals
>> Sign in here to start your access to the latest two volumes for 14 days

Full Article Figures & data References Citations Metrics

Reprints & Permissions Read this article

Abstract

The pathological effects of exposure to an electromagnetic field (EMF) during childhood and adolescence may be greater than those from exposure during adulthood. We investigated possible pathological changes in the cerebellum of adolescent rats exposed to 900 MHz EMF daily for 25 days. We used three groups of six 21-day-old male rats as follows: unexposed control group (Non-EG), sham-exposed group (Sham-EG) and an

EMF-exposed group (EMF-EG). EMF-EG rats were exposed to EMF in an EMF cage for 1 h daily from postnatal days 21 through 46. Sham-EG rats were placed in the EMF cage for 1 h daily, but were not subjected to EMF. No procedures were performed on the Non-EG rats. The cerebellums of all animals were removed on postnatal day 47, sectioned and stained with cresyl violet for histopathological and stereological analyses. We found significantly fewer Purkinje cells in the EMF-EG group than in the Non-EG and Sham-EG groups. Histopathological evaluation revealed alteration of normal Purkinje cell arrangement and pathological changes including intense staining of neuron cytoplasm in the EMF-EG group. We found that exposure to continuous 900 MHz EMF for 1 h/day during adolescence can disrupt cerebellar morphology and reduce the number of Purkinje cells in adolescent rats.

Key words: adolescence cell phone cerebellum electromagnetic field Purkinje cells rat

Previous article **View** issue table of contents **Next** article

Declaration of interest: The authors report no conflicts of interest. The authors alone are responsible for the content and writing of this paper.

Share

Related research

People also read Recommended articles Cited by 10

A review of curcumin as a biological stain and as a self-visualizing pharmaceutical agent

M Hope-Roberts et al.
Biotechnic & Histochemistry
Published online: 16 May 2017